

SUBCHORIONIC HEMATOMA'S INFLUENCE ON THE OUTCOME OF PREGNANCIES IN INDIVIDUALS FACING THREATENED ABORTION

Rozikova Dildora Kodirovna

Bukhara State Medical Institute, Bukhara, Uzbekistan

Introduction

Vaginal bleeding is a common complication during the first trimester of pregnancy, with an incidence ranging from 16% to 25% [1]. When there is intrauterine bleeding without cervical dilatation and tenderness in the early stages of pregnancy, it is referred to as a threatened abortion. Typically, this condition does not involve pain or excessive bleeding but can lead to maternal anxiety and potential adverse outcomes for both the mother and the fetus [2,4]. Placental dysfunction is one suggested mechanism for threatened abortion and has also been associated with various later complications, including preeclampsia, preterm labor, preterm birth, placental abruption, placenta previa, intrauterine growth restriction, and perinatal mortality [2, 3]. Inadequate angiogenesis is similarly linked to early pregnancy losses, and maternal serum AFP and β -hCG are proposed markers for angiogenesis in the first trimester [5]. Alongside these markers, chronic inflammation of the decidua may also underlie early pregnancy bleeding.

Ultrasound examinations frequently reveal intrauterine hemorrhages, especially in patients experiencing clinical bleeding in early pregnancy, with an incidence reported between 4% and 22% [6]. Subchorionic hematomas (SCHs) typically appear as hypoechoic or anechoic crescent-shaped areas on ultrasound. Although the exact cause is uncertain, they are believed to result from the partial detachment of the chorionic membranes from the uterine wall [7]. Possible predisposing factors include uterine malformations, a history of recurrent pregnancy loss, and infections [8,10]. The clinical significance of SCH remains a subject of debate [11,14], and it is unclear whether these hemorrhages directly lead to abortion. However, according to recent meta-analysis results, the presence of SCH increases the risk of both early and late pregnancy loss by a factor of two [15]. It is suggested that SCH presence elevates the risk of adverse obstetric outcomes, with fetal outcomes being influenced by factors such as the size of the hematoma, maternal age, and gestational age [16, 17].

The objective of this retrospective cohort study was to evaluate the impact of ultrasonographically identified SCH on pregnancy outcomes in patients experiencing vaginal bleeding during the first half of pregnancy.

Material and Methods

In this retrospective study, conducted between January 2021 and September 2023 at a university-based maternity hospital, 242 patients diagnosed with threatened

abortion due to painless vaginal bleeding were enrolled. The inclusion criteria were as follows: hospitalization due to threatened abortion, having a singleton pregnancy, gestational age less than 20 weeks, and continuous follow-up at the clinic until the end of the pregnancy. Exclusion criteria comprised a diagnosis of incipient abortion, absence of fetal cardiac activity, gestational age equal to or greater than 20 weeks, multiple pregnancies, and a history of recurrent pregnancy loss.

The study group encompassed 44 patients in whom subchorionic hematomas (SCHs) were detected through ultrasonography, while the control group included 198 patients without SCHs. All patients, regardless of group, were hospitalized for a minimum of 3 days after the initial episode of vaginal bleeding. Prophylactic progesterone treatment was administered to all patients, either orally. After being discharged from the hospital, patients followed standard antenatal care protocols. The study aimed to compare pregnancy outcomes between the two groups. For patients who experienced miscarriages, data on gestational age at the time of miscarriage and the duration between the initial bleeding episode and the miscarriage were recorded. In cases where pregnancies resulted in live births, comparisons were made regarding gestational age at delivery, birth weight, rates of preterm delivery, and cesarean section rates between the study and control groups.

Discussion

The results obtained from the present study revealed that the presence of SCH in patients with threatened abortion is an important factor for the continuation of pregnancy. The presence of SCH in patients with threatened abortion increases the risk of miscarriage. However, it does not affect the gestational age at miscarriage or the duration between the first bleeding and miscarriage. In patients whose pregnancies resulted in delivery, gestational age at labor, birth weight, preterm delivery, and cesarean section rates were not affected by the presence of SCH. Previously, several studies have investigated the effects of SCH on pregnancy outcomes. 238 patients with ultrasonographically detected SCH in a retrospective casecontrol study and reported a significant association between SCH and miscarriage and preterm delivery rates. They also reported increasing pregnancy loss rates with increasing SCH size. However, in the aforementioned studies, SCHs were defined during the routine first- or second-trimester ultrasonography, and not all patients with an SCH had threatened abortion.

Vaginal bleeding occurs in 25% of pregnancies in the first 20 weeks, and half of these result in miscarriage. Hence, it is important to identify the risk factors of threatened abortion and the factors that can affect the outcome. In a retrospective cohort trial, Ben-Haroush et al. assessed 2556 pregnant patients who were admitted with vaginal bleeding during the first 20 gestational weeks. The incidence of SCH was 9%. They reported that gestational age at diagnosis, size of SCH, and duration of bleeding did not affect the pregnancy outcome. They also reported significantly decreased

miscarriage rates with bed rest. However, Bennett et al. reported that increasing SCH size increases the risk of miscarriage. In a prospective cohort study, Pedersen and Mantoni followed up 342 pregnancies with vaginal bleeding between 9 to 20 gestational weeks, in which 18% had SCH. They found no association between the presence of SCH and miscarriage or preterm delivery risks. In another retrospective case-control study, Johns et al. reported that first-trimester vaginal bleedings were associated with adverse pregnancy outcomes, but the presence of SCH had no effect on the prognosis. According to the results of a recent meta-analysis evaluating 1735 patients with SCH from 7 studies, the presence of SCH increases the risks of early and late pregnancy loss, miscarriage, and preterm premature rupture of membranes.

In a prospective study from Uzbekistan, the size of the SCH was suggested to be the primary risk factor for miscarriage in patients with first-trimester vaginal bleeding. The first-trimester bleedings were associated with preterm delivery and low birth weight. However, we found no relationship between the prognosis and presence or size of the SCH.

The underlying mechanism of how SCH causes adverse pregnancy outcomes is still controversial. One of the possible mechanisms is the premature perfusion of the intervillous space, as occurs with subchorionic hemorrhage, before the development of placental adaptations to cope with oxidative stress [15]. Another possible mechanism might be the underlying cause of the subchorionic bleeding and secondary mechanical effects of the hematoma. Shallow trophoblast invasion and impaired angiogenesis with resultant friable blood vessels may predispose one to subchorionic hemorrhage, as well as adverse outcomes [10]. The presence of a hematoma, especially in a retroplacental location, may create an area of weakness, where further separation of the placenta from the uterine wall may occur, resulting in placental abruption [14]. Our results support the estimated mechanical effect of SCH that can cause miscarriage. The presence of an SCH and detachment of the gestational sac from the endometrium may result in miscarriage. However, if the gestational sac survives, reattachment to the endometrial wall might be enough for further progression of the pregnancy without any other adverse effects.

In conclusion, The presence of subchorionic hematoma (SCH) in individuals facing threatened abortion in the early stages of pregnancy elevates the likelihood of miscarriage. Yet, it remains uncertain whether the presence of SCH intensifies the risk of unfavorable pregnancy outcomes in continuing pregnancies, as the majority of past research has been retrospective. Substantial prospective randomized studies are needed to establish the actual impact of SCH on the prognosis of ongoing pregnancies.

References

1. Ikhtiyarova G.A., Iroda Tosheva, Nargiza Nasrullayeva. Causes of fetal loss syndrome at different gestation times //Asian Journal of Research. - 2017. - № 3 (3). - P. 32 - 41.
2. Rozikova D.K. The effect of subchorionic hematoma on pregnancy outcome in patients with threatened abortion (2023) Best journal of innovation in science, research and development Vol 2, issue: 10
3. Ikhtiyarova, G. A., Özcan, A. S., Gök, Ö., & Özcan, A. (2012). Characterization of natural-and organo-bentonite by XRD, SEM, FT-IR and thermal analysis techniques and its adsorption behaviour in aqueous solutions. *Clay Minerals*, 47(1), 31-44.
4. Ikhtiyarova, G. A., Tosheva, I. I., & Narzulloeva, N. S. (2017). Causes of fetal loss syndrome at different gestation times. *Asian Journal of Research*, 3(3).
5. Ikhtiyarova, G. A., Tosheva, I. I., Aslonova, M. J., & Dustova, N. K. (2020). Prenatal rupture of amnion membranes as A risk of development of obstetrics pathologies. *European Journal of Molecular and Clinical Medicine*, 7(7), 530-535. Retrieved from www.scopus.com
6. Ikhtiyarova, G. A., Tosheva, I. I., Aslonova, M. J., & Dustova, N. K. (2020). Prenatal rupture of amnion membranes as A risk of development of obstetrics pathologies. *European Journal of Molecular and Clinical Medicine*, 7(7), 530-535.
7. Rozikova D. K. (2023) The pattern of reproductive loses among women in Uzbekistan's population American kournal of pediatric medicine and health sciences Vol. 01 Issue 08, 2023 ISSN(E):2993-2149
8. Khashchenko E.P., Uvarova E.V. Modern approaches to the diagnosis and management of patients with polycystic ovary syndrome in adolescence // Obstetrics and gynecology. - 2015. - No. 5. - S. 5-9.
9. Kim M. W., Hong S. C., Choi J. S. Han J. Y. et al. Homocysteine, folate and pregnancy outcomes // J. Obstet. Gynaecol. 2012. Vol. 32. N 6. P. 520–524.
10. Kurbanova Z.Sh., Ikhtiyarova G.A., Rozikova D.K. - Anovulatory infertility and polycystic ovary syndrome // Tibbiyotdayangi kun 2 (30/2) 2020 pp. 159-165
11. Kurbanova Z.Sh., Ikhtiyarova G.A., Rozikova D.K. - Causes and methods of diagnosing endocrine infertility and the role of vitamin D in its correction // Tibbiyotdayangi kun 2 (30/2) 2020 pp. 34-38
12. Mavlyanova N.N, Ixtiyarova G.I, Tosheva I.I, et al. The State of the Cytokine Status in Pregnant Women with Fetal Growth Retardation. *J Med - ClinRes&Rev*. 2020; 4(6): 1-4.
13. Mavlyanova, N. N., Ixtiyarova, G. I., Tosheva, I. I., Aslonova, M. Z., & Narzullaeva, N. S. (2020). The State of the Cytokine Status in Pregnant Women with Fetal Growth Retardation. *J Med-Clin Res & Rev*, 4(6), 1-4.
14. Oripova F.Sh, Ikhtiyarova G. A., Kurbanova Z.Sh. Reproductive rehabilitation of women with chronic inflammatory diseases of the genitals.// Journal of biomedicine and practice special issue 2. - 2020. - P. 56 - 59.
15. Ихтиярова Г.А., Розикова Д. К. (2023). Микробиологические изменения у женщин с репродуктивными потерями. *Finland International Scientific Journal of Education, Social Science & Humanities*, 1002-1008
16. Mukhiddinovna, I. M. (2022). Effects of chronic consumption of energy drinks on liver and kidney of experimental rats. *International Journal of Philosophical Studies and Social Sciences*, 2(4), 6-11.

17. Mukhiddinova, I. M. (2022). EFFECTS OF CHRONIC CONSUMPTION OF ENERGY DRINKS ON LIVER AND KIDNEY OF EXPERIMENTAL RATS. *International Journal of Philosophical Studies and Social Sciences*, 2(4), 6-11.
18. Muxiddinova, I. M. (2022). Impact of energy drinks and their combination with alcohol to the rats metabolism. *Gospodarka i Innowacje.*, 22, 544-549.
19. Muxiddinova, I. M. (2022). IMPACT OF ENERGY DRINKS AND THEIR COMBINATION WITH ALCOHOL TO THE RATS METOBOLISM. *Gospodarka i Innowacje.*, 22, 544-549.
20. Muxiddinova, I. M. (2022). Effects of Energy Drinks on Biochemical and Sperm Parameters in Albino Rats. *Central Asian Journal of Medical and Natural Science*, 3(3), 126-131.
21. Muxiddinova, I. M. (2022). Demage of Energy Drinks on the Spermatogenesis of Male Rat's. *Research Journal of Trauma and Disability Studies*, 1(9), 111-118.
22. Muxiddinova, I. M. (2022). Effects of Energy Drinks on Biochemical and Sperm Parameters in Albino Rats. *Central Asian Journal of Medical and Natural Science*, 3(3), 126-131.
23. Muxiddinova, I. M. (2022). Impact of energy drinks and their combination with alcohol to the rats metabolism. *Gospodarka i Innowacje.*, 22, 544-549.
24. Muxiddinova, I. M. (2022). Ameliorative effect of Omega-3 on energy drinks-induced pancreatic toxicity in adult male albino rats. *International Journal of Health Systems and Medical Sciences*, 1(5), 13-18.
25. Muxiddinova, I. M., & Sobirova, A. Z. (2022). Pregnancy with Twins with Preeclampsia. *Central Asian Journal of Literature, Philosophy and Culture*, 3(11), 212-221.
26. Muxiddinova, I. M., & Sobirova, A. Z. (2022). Anemia Iron Deficiency in Pregnancy. *Central Asian Journal of Literature, Philosophy and Culture*, 3(11), 191-199.
27. Mukhiddinova, I. M. (2022). ENERGY DRINKS MAY AFFECT THE OVARIAN RESERVE AND SERUM ANTI-MULLERIAN HORMONE LEVELS IN A RAT MODEL. *BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI*, 2(12), 626-632.
28. Mukhiddinova, I. M. (2023). High Caffeine Exposure Increases Ovarian Estradiol Production in Immature Rats. *JOURNAL OF HEALTHCARE AND LIFE-SCIENCE RESEARCH*, 2(3), 8-11.
29. Mukhiddinova, I. M. (2023). Energy Fluids May Affect the Ovarian Reserve and Serum Anti-Mullerian Hormone Level. *Scholastic: Journal of Natural and Medical Education*, 2(5), 358-364.
30. Mukhiddinova, I. M. (2022). ENERGY DRINKS MAY AFFECT THE OVARIAN RESERVE AND SERUM ANTI-MULLERIAN HORMONE LEVELS IN A RAT MODEL. *BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI*, 2(12), 626-632.
31. Ismatova, M. M. (2023). Energy Drinks May Affect the Ovarium. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(8), 34-38.
32. Suratova, S. S., & Muxiddinova, I. M. (2023). Genetic Polymorphisms in Interleukin-1 β (Rs1143634) and Interleukin-8 (Rs4073) Are Associated With Survival after Resection of Intrahepatic Cholangiocarcinoma. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(8), 39-46.
33. Ismatova, M. M. (2023). Energy Drinks May Affect the Ovarium. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(8), 34-38.
34. Mukhiddinova, I. M. (2022). ENERGY DRINKS MAY AFFECT THE OVARIAN RESERVE AND SERUM ANTI-MULLERIAN HORMONE LEVELS IN A RAT

- MODEL. *BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI*, 2(12), 626-632.
35. Mukhiddinova, I. M. (2023). Energy Fluids May Affect the Ovarian Reserve and Serum Anti-Mullerian Hormone Level. *Scholastic: Journal of Natural and Medical Education*, 2(5), 358-364.
36. Muxiddinova, I. M. (2024). GENETIC POLYMORPHISMS IN INTERLEUKIN-1B (RS1143634) AND INTERLEUKIN-8 (RS4073) ARE ASSOCIATED WITH SURVIVAL AFTER RESECTION OF INTRANEPATIC CHOLANGIOCARCINOMA. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(6), 101-115.
37. Исмадова, М. М. (2024). ПРОГНОЗИРОВАНИЕ ВЕГЕТАТИВНЫХ НАРУШЕНИЙ У БЕРЕМЕННЫХ ПЕРЕНЕСШИХ COVID-19. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(6), 161-174.
38. Исмадова, М. М. (2024). ПРОГНОЗИРОВАНИЕ ДЕПРЕССИВНЫХ НАРУШЕНИЙ У БЕРЕМЕННЫХ И РОДИЛЬНИЦ ПОСЛЕ КОРОНАВИРУСНОЙ ИНФЕКЦИИ. *Journal of new century innovations*, 46(1), 140-151.
39. Muxiddinova, I. M. (2024). GENETIC POLYMORPHISMS IN INTERLEUKIN-1B (RS1143634) AND INTERLEUKIN-8 (RS4073) ARE ASSOCIATED WITH SURVIVAL AFTER RESECTION OF INTRANEPATIC CHOLANGIOCARCINOMA. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(6), 101-115.
40. Muxiddinova, I. M. (2024). GENETIC POLYMORPHISMS IN INTERLEUKIN-1B (RS1143634) AND INTERLEUKIN-8 (RS4073) ARE ASSOCIATED WITH SURVIVAL AFTER RESECTION OF INTRANEPATIC CHOLANGIOCARCINOMA. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(6), 101-115.
41. Исмадова, М. М. (2024). ПОСЛЕРОДОВЫХ ОСЛОЖНЕНИЙ. *Journal of new century innovations*, 46(1), 152-159.
42. Исмадова, М. М. (2024). ПОСЛЕРОДОВЫХ ОСЛОЖНЕНИЙ. *Journal of new century innovations*, 46(1), 152-159.
43. Исмадова, М. М. (2024). ВЕГЕТАТИВНЫХ НАРУШЕНИЙ У БЕРЕМЕННЫХ ПЕРЕНЕСШИХ COVID-19. *Journal of new century innovations*, 46(1), 129-132.
44. Ismatova, M. M. (2024). PROBLEMS OF LATE POSTPARTUM COMPLICATIONS AND ITS CORRECTION. *Journal of new century innovations*, 46(1), 160-167.
45. Ихтиярова, Г. А., & Розикова, Д. К. (2023). МИКРОБИОЛОГИЧЕСКИЕ ИЗМЕНЕНИЯ У БЕРЕМЕННЫХ С РЕПРОДУКТИВНЫМИ ПОТЕРЯМИ В АНАМНЕЗЕ. *Finland International Scientific Journal of Education, Social Science & Humanities*, 11(4), 1002-1008.
46. Розикова, Д. К., & Ихтиярова, Г. А. (2023). THE STRUCTURE OF REPRODUCTIVE LOSSES IN UZBEK WOMEN. *ЖУРНАЛ РЕПРОДУКТИВНОГО ЗДОРОВЬЯ И УРОНЕФРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ*, 4(4).
47. Rozikova, D. K. (2023). THE IMPACT OF SUBCHORIONIC HEMATOMA ON THE FINAL RESULT OF PREGNANCIES IN INDIVIDUALS EXPERIENCING THREATENED ABORTION. *GOLDEN BRAIN*, 1(28), 57-62.
48. Rozikova Dildora Kodirovna. (2023). The Pattern of Reproductive Losese among Women in Uzbekistan's Population. *American Journal of Pediatric Medicine and Health Sciences* (2993-2149), 1(8), 52-60.
49. Kodirovna, R. D. (2023). The Effects of Subchorionic Hematoma on Pregnancy Outcome in Patients with Threatened Abortion. *Best Journal of Innovation in Science, Research and Development*, 2(10), 121-124.

50. Ikhtiyarova, G. A., Dustova, N. K., & Qayumova, G. (2017). Diagnostic characteristics of pregnancy in women with antenatal fetal death. *European Journal of Research*, (5), 5.
51. Kayumova, G. M., & Nutfilloyevich, K. K. (2023). CAUSE OF PERINATAL LOSS WITH PREMATURE RUPTURE OF AMNIOTIC FLUID IN WOMEN WITH ANEMIA. *AMALIY VA TIBBIYOT FANLARI ILMIIY JURNALI*, 2(11), 131-136.
52. Kayumova, G. M., & Dustova, N. K. (2023). Significance of the femoflor test in assessing the state of vaginal microbiocenosis in preterm vaginal discharge. Problems and scientific solutions. In *International conference: problems and scientific solutions. Abstracts of viii international scientific and practical conference* (Vol. 2, No. 2, pp. 150-153).
53. Каюмова, Г. М., Мухторова, Ю. М., & Хамроев, Х. Н. (2022). Определить особенности течения беременности и родов при дородовом излитии околоплодных вод. *Scientific and innovative therapy. Научный журнал по научный и инновационный терапии*, 58-59.
54. Kayumova, G. M., & Dustova, N. K. (2023). ASSESSMENT OF THE STATE OF THE GENITAL TRACT MICROBIOCENOSIS IN PREGNANT WOMEN WITH PREMATURE RUPTURE OF THE MEMBRANES USING THE FEMOFLOR TEST. *Modern Scientific Research International Scientific Journal*, 1(1), 70-72.
55. Valeryevna, S. L., Mukhtorovna, K. G., & Kobyllovna, E. S. (2019). Premature Birth In A Modern Aspect. *International Journal of Bio-Science and Bio-Technology*, 11(10), 31-37.
56. Саркисова, Л. В., Каюмова, Г. М., & Умидова, Н. Н. (2018). Морфологические изменения фетоплацентарного комплекса при герпетической инфекции. *Тиббиётда янги кун*, 188-191.
57. Каюмова, Г. М., Саркисова, Л. В., & Умидова, Н. Н. (2018). Современные взгляды на проблему преждевременных родов. *Тиббиётда янги кун*, 183-185.
58. Каюмова, Г. М., Хамроев, Х. Н., & Ихтиярова, Г. А. (2021). Причины риска развития преждевременных родов в период пандемии организм и среда жизни к 207-летию со дня рождения Карла Францевича Рулье: сборник материалов IV-ой Международной научнопрактической конференции (Кемерово, 26 февраля 2021 г.). ISBN 978-5-8151-0158-6.139-148.
59. Саркисова, Л. В., Каюмова, Г. М., & Бафаева, Н. Т. (2019). Причины преждевременных родов и пути их решения. *Биология ва тиббиёт муаммолари*, 115(4), 2.
60. Kayumova, G. M., & Dustova, N. K. (2023). Significance of the femoflor test in assessing the state of vaginal microbiocenosis in preterm vaginal discharge. Problems and scientific solutions. In *International conference: problems and scientific solutions. Abstracts of viii international scientific and practical conference* (Vol. 2, No. 2, pp. 150-153).
61. KAYUMOVA, G., & DUSTOVA, N. (2023). Features of the hormonal background with premature surge of amniotic fluid. *Of the international scientific and practical conference of young scientists «Science and youth: conference on the quality of medical care and health literacy» Ministry of healthcare of the republic of kazakhstan kazakhstan's medical university «KSPH»*. ISBN 978-601-305-519-0.29-30.
62. Каюмова, Г. М. НҚ Дўстова.(2023). Muddatdan oldin qog'onoq suvining ketishida xavf omillarning ta'sirini baholash. *Журнал гуманитарных и естественных наук*, 2(07), 11-18.
63. Каюмова, Г. М., & Мухторова, Ю. М. (2022). Пороговые значения антител к эстрадиолу, прогестерону и бензо [a] пирену как факторы риска преждевременного излития околоплодных вод при недоношенной беременности. *Scientific and innovative therapy. Научный журнал по научный и инновационный терапии*, 59-60.

64. Каюмова, Г. М., Мухторова, Ю. М., & Хамроев, Х. Н. (2022). Причина преждевременных родов. *Scientific and innovative therapy. Научный журнал по научный и инновационный терапии*, 57-58.
65. Sarkisova, L. V., & Kayumova, G. M. (2019). Exodus of premature birth. *Тиббиётда янги кун*, 1(25), 155-159.
66. Саркисова, Л. В., & Каюмова, Г. М. (2018). Перинатальный риск и исход преждевременных родов. *Проблемы медицины и биологии*, 169-175.
67. Каюмова, Г. М., Саркисова, Л. В., & Рахматуллаева, М. М. (2018). Особенности состояния плаценты при преждевременных родах. In *Республиканской научно практической конференции «Актуальные вопросы охраны здоровья матери и ребенка, достижения и перспективы»* (pp. 57-59).
68. Каюмова, Г. М., Саркисова, Л. В., & Саъдуллаева, Л. Э. (2018). Показатели центральной гемодинамики и маточно-фетоплацентарного кровотока при недонашивании беременности. In *Республиканской научно практической конференции «Актуальные вопросы охраны здоровья матери и ребенка, достижения и перспективы»* (pp. 56-57).
69. Саркисова, Л., Каюмова, Г., & Рузиева, Д. (2019). Современные тренды преждевременных родов. *Журнал вестник врача*, 1(4), 110-114.
70. Каюмова, Г. М., & Ихтиярова, Г. А. (2021). Причина перинатальных потер при преждевременных родов у женщин с анемией.(2021). In *Материалы республиканской научно-практической онлайн конференции.«Актуальные проблемы современной медицины в условиях эпидемии»* (pp. 76-7).
71. Kayumova, G. M., Khamroev, X. N., & Ixtiyarova, G. A. (2021). Morphological features of placental changes in preterm labor. *Тиббиётда янги кун*, 3(35/1), 104-107.
72. Khamroyev XN, Q. G. (2021). Improving the results of treatment of choledocholithiasis in liver diseases.
73. Kayumova, G. M. (2023). TO DETERMINE THE FEATURES OF THE COURSE OF PREGNANCY AND CHILDBIRTH IN WOMEN WITH PRENATAL RUPTURE OF AMNIOTIC FLUID. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(11), 137-144.
74. Kayumova, G. M. (2023). To Determine the Features Of Pregnancy and Children During Antenature Ruption Of Ambient Fluid. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 66-72.
75. Kayumova, G. M. (2023). Features of the Hormonal Background During Premature Relation of Ambitionnal Fluid. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 73-79.
76. Kayumova, G. M. (2023). The Significance Of Anti-Esterogen And Progesterone Antibodies As A Risk Factor In Premature Rupture Of Amniotic Fluid. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 58-65.
77. Каюмова, Г. М. (2024). ПЕРИНАТАЛЬНЫЕ ИСХОДЫ ПРИ ДОРОДОВОМ РАЗРЫВЕ ПЛОДНЫХ ОБОЛОЧЕК. *Journal of new century innovations*, 46(1), 242-251.
78. Каюмова, Г. М. (2024). ОПРЕДЕЛИТЬ ФАКТОРЫ РИСКА ПРЕЖДЕВРЕМЕННЫХ РОДОВ. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 38(6), 228-235.
79. Каюмова, Г. М. (2024). ОСОБЕННОСТИ КАЧЕСТВЕННОГО СОСТАВА МИКРОБИОТА ВЛАГАЛИЩА ПРИ АКТИВНО-ВЫЖИДАТЕЛЬНОЙ ТАКТИКЕ ВЕДЕНИЯ БЕРЕМЕННЫХ С ПРЕЖДЕВРЕМЕННЫМ ИЗЛИТИЕМ ОКОЛОПЛОДНЫХ ВОД. *Journal of new century innovations*, 46(1), 231-241.

80. Каюмова, Г. М. (2024). ИССЛЕДОВАНИЕ МИКРОБИОТА ВЛАГАЛИЩА ПРИ ДОРОДОВОМ ИЗЛИТИИ ОКОЛОПЛОДНЫХ ВОД. *Journal of new century innovations*, 46(1), 213-221.
81. Каюмова, Г. М. (2024). ИССЛЕДОВАНИЕ ПОКАЗАТЕЛЕЙ КРОВИ У БЕРЕМЕННЫХ С ПРЕЖДЕВРЕМЕННЫМИ ОКОЛОПЛОДНЫМИ ВОДАМИ. *Journal of new century innovations*, 46(1), 222-230.
82. Kayumova, G. M. (2024). ANTIBACTERIAL THERAPY FOR PRETERMARY AND ANTENATURE RURUSION OF AMBITIONAL FLUID. *Journal of new century innovations*, 46(1), 252-262.
83. Уроков, Ш. Т., & Хамроев, Х. Н. (2019). Influe of diffusion diseases of the liver on the current and forecst of obstructive jaundice. *Тиббиётда янги кун*, 1, 30.
84. TESHAEV, S. J., TUHSANOVA, N. E., & HAMRAEV, K. N. (2020). Influence of environmental factors on the morphometric parameters of the small intestine of rats in postnatal ontogenesis. *International Journal of Pharmaceutical Research (09752366)*, 12(3).
85. Хамроев, Х. Н. (2022). Toxic liver damage in acute phase of ethanol intoxication and its experimental correction with chelate zinc compound. *European journal of modern medicine and practice*, 2, 2.
86. Khamroev, B. S. (2022). RESULTS OF TREATMENT OF PATIENTS WITH BLEEDING OF THE STOMACH AND 12 DUO FROM NON-STEROIDAL ANTI-INFLAMMATORY DRUGS-INDUCED OENP. *Journal of Pharmaceutical Negative Results*, 1901-1910.
87. Nutfilloyevich, K. K. (2023). STUDY OF NORMAL MORPHOMETRIC PARAMETERS OF THE LIVER. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(8), 302-305.
88. Nutfilloyevich, K. K. (2024). NORMAL MORPHOMETRIC PARAMETERS OF THE LIVER OF LABORATORY RATS. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(3), 104-113.
89. Nutfilloeovich, K. K., & Akhrorovna, K. D. (2024). MORPHOLOGICAL CHANGES IN THE LIVER IN NORMAL AND CHRONIC ALCOHOL POISONING. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(3), 77-85.
90. Kayumova, G. M., & Hamroyev, X. N. (2023). SIGNIFICANCE OF THE FEMOFLOR TEST IN ASSESSING THE STATE OF VAGINAL MICROBIOCENOSIS IN PRETERM VAGINAL DISCHARGE. *International Journal of Medical Sciences And Clinical Research*, 3(02), 58-63.
91. Хамроев, Х. Н., & Тухсанова, Н. Э. (2022). НОВЫЙ ДЕНЬ В МЕДИЦИНЕ. *НОВЫЙ ДЕНЬ В МЕДИЦИНЕ Учредители: Бухарский государственный медицинский институт, ООО "Новый день в медицине"*, (1), 233-239.
92. Хамроев, Х. Н. (2024). Провести оценку морфологических изменений печени в норме и особенностей характера ее изменений при хронической алкогольной интоксикации. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(3), 95-3.
93. Хамроев, Х. Н., & Туксанова, Н. Э. (2021). Characteristic of morphometric parameters of internal organs in experimental chronic alcoholism. *Тиббиётда янги кун*, 2, 34.
94. Хамроев, Х. Н., Хасанова, Д. А., Ганжиев, Ф. Х., & Мусоев, Т. Я. (2023). Шошилинич тиббий ёрдам ташкил қилишининг долзарб муаммолари: Политравма ва ўткир юрак-қон томир касалликларида ёрдам кўрсатиш масалалари. *XVIII Республика илмий-амалий анжумани*, 12.

95. Хамроев, Х. Н., & Хасанова, Д. А. (2023). Жигар морфометрик кўрсаткичларининг меъёра ва экспериментал сурункали алкохолизмда қиёсий таснифи. *Медицинский журнал Узбекистана | Medical journal of Uzbekistan*, 2.
96. Khamroyev, X. N. (2022). TOXIC LIVER DAMAGE IN ACUTE PHASE OF ETHANOL INTOXICATION AND ITS EXPERIMENTAL CORRECTION WITH CHELATE ZINC COMPOUND. *European Journal of Modern Medicine and Practice*, 2(2), 12-16.
97. Xamroyev, X. N. (2022). The morphofunctional changes in internal organs during alcohol intoxication. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 2(2), 9-11.
98. Khamroyev, X. N. (2022). TOXIC LIVER DAMAGE IN ACUTE PHASE OF ETHANOL INTOXICATION AND ITS EXPERIMENTAL CORRECTION WITH CHELATE ZINC COMPOUND. *European Journal of Modern Medicine and Practice*, 2(2), 12-16.
99. Xamroyev, X. N. (2022). The morphofunctional changes in internal organs during alcohol intoxication. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 2(2), 9-11.
100. Латипов, И. И., & Хамроев, Х. Н. (2023). Улучшение Результат Диагностики Ультразвуковой Допплерографии Синдрома Хронической Абдоминальной Ишемии. *Central Asian Journal of Medical and Natural Science*, 4(4), 522-525.
101. Хамроев, Х. Н., & Уроков, Ш. Т. (2019). ВЛИЯНИЕ ДИФФУЗНЫХ ЗАБОЛЕВАНИЙ ПЕЧЕНИ НА ТЕЧЕНИЕ И ПРОГНОЗ МЕХАНИЧЕСКОЙ ЖЕЛТУХИ. *Новый день в медицине*, (3), 275-278.
102. Хамроев, Х. Н., & Ганжиев, Ф. Х. (2023). Динамика структурно-функциональных нарушение печени крыс при экспериментальном алкохолие циррозе. *Problems of modern surgery*, 6.
103. Уроков, Ш. Т., & Хамроев, Х. Н. (2018). Клинико-диагностические аспекты механической желтухи, сочетающейся с хроническими диффузными заболеваниями печени (обзор литературы). *Достижения науки и образования*, (12 (34)), 56-64.
104. Nutfilloevich, N. K., & Akhrorovna, K. D. (2023). COMPARATIVE CLASSIFICATION OF LIVER MORPHOMETRIC PARAMETERS IN THE LIVER AND IN EXPERIMENTAL CHRONIC ALCOHOLISM. *International Journal of Cognitive Neuroscience and Psychology*, 1(1), 23-29.