

CLINICAL AND MORPHOLOGICAL CHANGES IN ORAL ORGANS AND TISSUES IN WOMEN AFTER MENOPAUSE AND THEIR ANALYSIS

Chakkanov Fakhritdin Khusanovich

Assistant Department of orthopedic dentistry

Samarkand State Medical University

Samarkand, Uzbekistan.

Annotation. During menopause, irreversible changes in the hormonal function of the ovaries occur, the concentration of estrogens decreases.

It is during this period that a progressive increase in various diseases is noted [8]. The study of the dental status in menopausal women has proven the existence of a relationship between a decrease in estrogen concentration and a sharp increase in dental diseases [4]. Because the oral mucosa and salivary glands contain estrogen receptors, changes in hormonal levels can be seen directly in the mouth. Leimola-Virtanen et al confirmed the presence of estrogen mRNA and immunoreactive protein in the oral mucosa and salivary glands, which also confirms its biological role [11].

Thus, the climacteric period is characterized by a disorder of metabolism and function in the tissues of the oral cavity [6]. As a result, tooth loss, periodontal disease, increased bone resorption of the alveolar processes are observed.

Keywords: menopause, generalized periodontitis, postmenopausal period, diseases of the oral mucosa.

The purpose of the study: to study the clinical and microbiological manifestations and mechanisms of development of dental diseases in postmenopausal women.

Material and methods: we examined 40 postmenopausal women with various diseases of the oral cavity. The control group of patients were women who are not in the postmenopausal period with dental diseases.

Results and discussions: All participants of the study underwent a comprehensive examination: hygienic and periodontal indices (OHI-S hygiene index, PI periodontal index), pH of mixed saliva was measured, sialometry was performed according to the method of M.M. Pozharitskaya, studied the microflora of the oral cavity by PCR (polymerase chain reaction).

When collecting anamnesis, it was revealed that most often women of group I complained of bleeding gums (13.3%) and bad breath (6.7%). Group II patients complained of bleeding gums (66%), bad breath (43.9%), exposure of the roots of the teeth (15.3%), tooth mobility (16.1%), and 40% of women had complaints of dry

mouth. While 26.7% of postmenopausal women taking HRT complained of bleeding gums, 18.1% complained of dry mouth and bad breath.

According to the results of the survey, postmenopausal women taking HRT 2.3 times less complained of bleeding gums (27.7%) and dry mouth (17.1%) compared with women not taking HRT (63%; 40% respectively). 18.1% of women taking HRT noted bad breath, which is 2.5 times less than women not taking HRT (43.9%), but 3 times more than in the control group (5.8%). Postmenopausal women, regardless of whether they take HRT or not, were 3 times more likely to have a white coating on the tongue (18.1%, 15.3%, respectively) than women in the control group (6.7%). Burning sensation in the oral cavity was noted by 5.7% of women in group II and 3.9% in group III.

Attention was also paid to the TMJ, noting the symmetry of the face and movements of the mandible when opening the mouth, pain when moving the mandible, and the range of vertical and lateral movements of the mandible. In 22.6% of group I 49, in 40% of group II and 45.7% of group III women, a deviation was observed, manifested by a displacement of the lower jaw to the side at the beginning and a return to the midline in the middle of opening the mouth by 2 mm. None of the subjects had limited range of vertical and lateral movements. 2.9% - group I, 14.3% - group II and 11.5% - group III sometimes noted clicks during the movement of the lower jaw.

When examining the oral cavity, such non-carious lesions of the teeth as erosion were noted, which were detected in 2.8% of women in group II. Wedge-shaped defects were noted in 17.1% of group I, 54.3% of group II and 62.8% of group III women. Pathological tooth wear was detected in 17.1% of group II and 14.2% of group III women, but there were no significant differences between the groups ($p > 0.05$).

The clinical examination included the determination of the KPU index, the OHI-S hygiene index, the periodontal index PI, the measurement of the pH of the mixed saliva, sialometry, and the assessment of the microflora of the oral cavity by PCR (polymerase chain reaction).

In the study of the intensity of tooth decay by caries, it was found that in group I, the average value of the KPU index = 13.58 ± 6.35 (moderate level of caries intensity), in group II, KPU = 18.8 ± 6.17 (very high level of caries intensity), and in group III, the KPU index = 16.45 ± 6.2 (high level of caries intensity).

When examining the level of oral hygiene, it was found that the OHI-S hygiene index, which takes into account both the amount of plaque and tartar, in group I was 1.83 ± 1.1 , in group II - 2.89 ± 0.89 , in group III - 3.24 ± 1.6 . A statistically significant difference was found between groups I and II, groups II and III ($p < 0.05$), and between groups I and III there were no significant differences in relation to the values of group parameters ($p > 0.05$).

To determine the severity of periodontal disease, the periodontal index PI was used, which showed that its value in women who do not take HRT is 2.31 ± 1.29 , which corresponds to the average degree of periodontal pathology and is 2 times more than in women taking HRT and 5 times more than in the control group. In women taking HRT, the value of the periodontal index PI is 1.16 ± 0.75 , which corresponds to the initial and mild degree of periodontal pathology, and in the control group - 0.46 ± 0.56 .

An objective examination of patients showed increased saliva viscosity, combined with foaminess in 51.4% of women not taking HRT, and in 25.7% of women taking HRT.

Sialometry according to the method of M.M. Pozharitskaya allows you to determine the amount of mixed unstimulated saliva. The greatest hyposalivation in the oral cavity was observed in women not taking HRT, and the average for the group was 2.1 ± 0.28 ml at a rate of 4.1 ml.

In the study of the pH of mixed saliva in women in group I, the average value reached 7.04 ± 0.35 , in group II, $pH = 6.48 \pm 0.33$, and in group III, the average $pH = 6.94 \pm 0.36$.

Analysis of samples of the contents of periodontal pockets showed that *A. actinomycetemcomitans*, a gram-negative non-motile facultative anaerobic coccobacillus, was found in 22.9% of women in group I, 51.4% in group II and 37.1% in group III. It was found that *A. actinomycetemcomitans* occurs 1.4 times more often in women who do not take HRT compared to women taking it and 2.2 times more often than in the control group. At the same time, a significant difference ($p < 0.05$) is observed only between groups I and II.

The frequency of occurrence of *T. denticola*, which is able to organize associations with other bacteria, especially with *P. gingivalis* and *T. forsythia*, is 37.1% in group I, 54.3% in group II and 51.4% in group III, but no significant difference ($p > 0.05$) between groups was found. *T. forsythia*, which most strongly promotes cell apoptosis, was found in 51.4% in group I, in 80% in group II, and in 45.7% in group III. A significant difference was found between groups I and II, II and III. Periodontopathogen *P. gingivalis*, as the most aggressive bacterium, was found in 11.4% of women in group I, 77.1% in group II and 28.6% in group III, respectively. It was shown that *P. gingivalis* occurs 6.7 times more often in group II than in group I and 2.7 times more often than in group III.

P. intermedia was found in 11.3% of women in group I, in 74.3% in group II and in 25.7% in group III, respectively. *P. intermedia* is 6.6 times more common in group II than in group I and 2.9 times more common than in group III. At the same time, there is a significant difference between *P. gingivalis* and *P. intermedia* ($p < 0.05$) between groups I and II, II and III. There was no significant difference between the qualitative

composition of the microflora of periodontal pockets in groups I and III (control and the group of women taking HRT) ($p > 0.05$).

Most often, in women not taking HRT, *P. gingivalis*, *P. intermedia*, and *T. forsythia* were detected, in contrast to women taking HRT. *A. actinomycetemcomitans* and *T. denticola* as part of the subgingival microflora were found regardless of the use of HRT and were detected in a smaller number of patients.

The study also revealed Epstein-Barr virus in 11.4% of women in group I, 20% in group II and 17.1% in group III. Herpes simplex virus was found in 17.1% of the women of group I, 25.7% of group II and 28.6% of group III. *Candida albicans* was found in 5.7% of women of group I, 25.7% of group II and 20% of group III, and a significant difference was found between groups I and II, II and III ($p < 0.05$).

In all patients of the control group, in whom a periodontopathogen was detected, the amount was relatively low and did not exceed 105 units/ml.

Spearman's rank correlation method was used to identify the relationship between the degree of clinical indicators and the composition of the microflora of periodontal pockets.

The relationship between an increased number of *A. actinomycetemcomitans* in the contents of the periodontal pocket with a low level of oral hygiene according to the OHI-S index ($r=0.503027$) and with the severity of periodontitis ($r=0.505208$) was revealed. A negative correlation was observed with the pH of the mixed saliva ($r=-0.456773$).

T. denticola

No correlation was found between the amount of *T. denticola* in periodontal pockets and clinical parameters.

P. gingivalis

The content of *P. gingivalis* moderately correlated with the level of oral hygiene ($r=0.469771$). A negative relationship between the content of *P. gingivalis* and the pH of mixed saliva ($r=-0.491576$) was revealed.

T. forsythia

The amount of *T. forsythia* in the contents of periodontal pockets moderately correlated with the level of oral hygiene ($r=0.376309$), and to a greater extent with the severity of periodontitis ($r=0.552702$). A negative correlation was also found with the pH of the mixed saliva.

P. intermedia

The level of *P. intermedia* depended on the severity of periodontitis ($r=0.416681$).

According to the results of the study, a statistically significant relationship was revealed between oral hygiene and the severity of periodontitis with the quantitative composition of periodontopathogens of group II ($p < 0.05$).

Findings. Thus, the greater sensitivity of the above indices is associated with the peculiarities of the pathogenesis of menopause, namely with estrogen deficiency. The content of estrogen receptors on the oral mucosa plays a direct role in the development of dental diseases [9]. During menopause, the angioprotective effect of estrogens weakens, followed by damage to the structures of the hemomicrocirculatory bed. Also during this period, the production of osteoclasts increases, the production of osteoblasts decreases, the absorption of calcium in the intestine decreases, the lack of vitamin D, which leads to increased bone resorption.

Due to estrogen deficiency, there is a decrease in intestinal absorption of calcium in the body, which in turn leads to disturbances in the regulation of calcium-phosphate metabolism and increased calcium release not only into the blood serum, but also into saliva [6]. Therefore, a high concentration of calcium in the saliva of menopausal women can lead to faster plaque mineralization and, consequently, to an increase in stone formation, which has a direct impact on the progression of gingivitis and periodontitis [9;1]. Thus, the pathogenesis of menopause affects the state of all structures of the oral cavity.

Literature:

1. Alimdjanovich, R. J., & Astanovich, A. A. (2023). Development of dental care in uzbekistan using a conceptual approach to improve its quality. *Central Asian Journal of Medical and Natural Science*, 4(4), 536-543.
2. Alimdjanovich, R. J., & Astanovich, A. A. (2023). FUNDAMENTALS OF DENTAL CARE IN THE REPUBLIC OF UZBEKISTAN BASED ON THE DEVELOPMENT OF GENERAL MEDICAL PRACTICE. *Best Journal of Innovation in Science, Research and Development*, 2(8), 163-174.
3. Astanovich, A. D. A., Alimdjanovich, R. J., Abdujamilevich, S. A., & Bakhriiddnovich, T. A. (2021). The State of Periodontal Tissues in Athletes Engaged in Cyclic Sports. *Annals of the Romanian Society for Cell Biology*, 235-241.
4. Bustanovna, I. N. (2022). Assessment of clinical and morphological changes in the oral organs and tissues in post-menopause women. *Thematics Journal of Education*, 7(3).
5. Bustanovna, I. N., & Sharipovna, N. N. (2023). Research cases in women after menopause clinical and morphological changes in oral organs and their analysis. *Journal of biomedicine and practice*, 8(3).
6. Bustanovna, I. N. (2024). Complications Arising in the Oral Cavity after Polychemotherapy in Patients with Hemablastoses. *International Journal of Scientific Trends*, 3(3), 62-66.

7. Bustanovna, I. N. (2024). The Effectiveness of the Use of the Drug "Proroot MTA" in the Therapeutic and Surgical Treatment of Periodontitis. *International Journal of Scientific Trends*, 3(3), 72-75.
8. Bustanovna, I. N., & Abdusattor o'g, A. A. A. (2024). Analysis of Errors and Complications in the Use of Endocal Structures Used in Dentistry. *International Journal of Scientific Trends*, 3(3), 82-86.
9. Bustanovna, I. N. (2024). Morphological Changes in Oral Organs and Tissues in Women after Menopause and their Analysis. *International Journal of Scientific Trends*, 3(3), 87-93.
10. Bustanovna, P. I. N. (2024). Further Research the Features of the Use of Metal-Ceramic Structures in Anomalies of Development and Position of Teeth. *International Journal of Scientific Trends*, 3(3), 67-71.
11. Bustanovna, P. I. N. (2024). Research of the Structure of Somatic Pathology in Patients with Aphthous Stomatitis. *International Journal of Scientific Trends*, 3(3), 51-55.
12. Husanovich, C. F. (2024). RESEARCH ON THE ATTITUDE OF PEOPLE TO THE PREVENTION OF DENTAL DISEASES. *European International Journal of Multidisciplinary Research and Management Studies*, 4(02), 265-268.
13. Исламова Н., Чакконов Ф. Роль продуктов перекисного окисления липидов и противовоспалительных цитокинов крови в развитии заболеваний полости рта при гипотиреозе //Общество и инновации. – 2020. – Т. 1. – №. 1/s. – С. 577-582.
14. Исламова, Н. Б. (2024). Complications Arising in the Oral Cavity after Polychemotherapy in Patients with Hemablastosis. *International Journal of Scientific Trends*, 3(3), 76-81.
15. Исламова, Н. Б., & Чакконов, Ф. Х. (2021). Изменения в тканях и органах рта при эндокринных заболеваниях. In *Актуальные вопросы стоматологии* (pp. 320-326).
16. Islamova, N. B., & Chakkonov, F. K. (2021). Changes in the tissues and organs of the mouth in endocrine diseases. *Current Issues in Dentistry*, 320-326.
17. Islamova, N. B., & Sh, N. N. (2023, May). STUDY OF CHANGES IN PERIODONTAL DISEASES IN POSTMENOPAUSAL WOMEN. In *Conferences* (pp. 15-17).
18. Khusanovich, C. F. (2023). Improvement of the Orthopedic Method in Complex Treatment in Patients with Periodontal Diseases. *Progress Annals: Journal of Progressive Research*, 1(3), 27-30.
19. Khusanovich, C. F. (2022). CHANGES IN TISSUES AND ORGANS OF THE MOUTH WITH ENDOCRINE DISEASES. *Thematics Journal of Education*, 7(3).

20. Khusanovich, K. B. R. C. F. (2023). Types and applications of dental compliments. *Journal of Modern Educational Achievements*, 5(5), 95-99.
21. Khusanovich, C. F. (2023). Therapeutic and Surgical Methods Treatment of Localized Periodontitis. *Scholastic: Journal of Natural and Medical Education*, 2, 142-148.
22. Nazarova, N. S., & Islomova, N. B. (2022). postmenopauza davridagi ayollarda stomatologik kasalliklarining klinik va mikrobiologik ko'rsatmalari va mexanizmlari. *Журнал "Медицина и инновации"*, (2), 204-211.
23. Sulaymonova, Z. Z., & Islamova, N. B. (2023, May). TAKING IMPRESSIONS IN THE ORAL CAVITY AND THEIR REDUCTION. In *Conferences* (pp. 21-23).
24. Sharipovna, N. N., & Bustanovna, I. N. (2022). Assessment of clinical and morphological changes in the oral organs and tissues in post-menopause women. *Frontline medical sciences and pharmaceutical journal*, 2(05), 60-67.
25. ЧАККОНОВ, Ф. Х. (2021). ЯТРОГЕННЫЕ ОШИБКИ В СТОМАТОЛОГИИ И ИХ ПРИЧИНЫ. In *Актуальные вопросы стоматологии* (pp. 925-930).
26. ЧАККОНОВ, Ф., САМАДОВ, Ш., & ИСЛАМОВА, Н. (2022). ENDOKANAL PIN-KONSTRUKSIYALARNI ISHLATISHDA ASORATLAR VA XATOLAR TAHLILI. *ЖУРНАЛ БИОМЕДИЦИНЫ И ПРАКТИКИ*, 7(1).
27. Xusanovich, C. F., Orzimurod, T., Maruf, U., & Ollomurod, X. (2023). PROSTHETICS A COMPLETE REMOVABLE PROSTHESIS BASED ON IMPLANTS. *European International Journal of Multidisciplinary Research and Management Studies*, 3(11), 122-126.
28. Xusanovich, C. F., Sunnat, R., & Sherali, X. (2024). CLASP PROSTHESES–TECHNOLOGY IMPROVEMENT. *European International Journal of Multidisciplinary Research and Management Studies*, 4(03), 152-156.