

UDK: 618.396: 616-09-067.15

DEVELOPMENT OF PREMATURE BIRTH DUE TO VITAMIN D DEFICIENCY

Sharipova N.M.

Bukhara State Medical Institute, Bukhara, Uzbekistan

Relevance. The problem of preterm birth (PTB) continues to worry not only the obstetric community, but also the state as a whole, because PTB is an important socio-economic problem, poses a threat to the life of mother and child, leads to early perinatal mortality and (or) disability of premature infants. In pregnant women, syncytiotrophoblast is one of the sources of calcitriol production. During pregnancy, closer to 12 weeks, the concentration of 1,25(OH)₂D more than doubles, which confirms the important role of vitamin D in supporting gestation and controlling fetal development processes. It regulates decidualization, implantation, and expression of placental lactogen. In addition, vitamin D affects the secretion of progesterone and estriol, human chorionic gonadotropin (hCG), calcium absorption by the placenta and placental immunomodulation. Vitamin D plays an important role in maintaining the mother-placenta-fetus system, including implantation, placental formation, embryogenesis, intra- and postnatal periods.

Key words: vitamin D deficiency, 25-hydroxyvitamin D, general risk groups, pregnancy, preterm birth.

РАЗВИТИЕ ПРЕЖДЕВРЕМЕННЫХ РОДОВ НА ФОНЕ ДЕФИЦИТА ВИТАМИНА D

Шарипова Н.М.

Бухарский государственный медицинский институт, Бухара, Узбекистан.

Актуальность. Проблема преждевременных родов (ПР) не перестает волновать не только акушерское сообщество, но и государство в целом, потому что ПР являются важной социально-экономической проблемой, несут угрозу жизни матери и ребенка, приводят к ранней перинатальной смертности и (или) к инвалидизации недоношенных детей. У беременных женщин синцитиотрофобласт является одним из источников продукции кальцитриола. Во время беременности, ближе к 12 нед, концентрация 1,25(OH)₂D увеличивается более чем вдвое, что подтверждает важную роль витамина D в поддержке гестации и контроле процессов развития плода. Он регулирует децидуализацию, имплантацию, экспрессию плацентарного лактогена. Кроме того, витамин D влияет на секрецию прогестерона и эстриола, хорионического

гонадотропина человека (ХГЧ), на поглощение кальция плацентой и плацентарную иммуномодуляцию. Витамин D играет важную роль в обеспечении системы мать-плацента-плод, включая имплантацию, формирование плаценты, эмбриогенез, интра-и постнатальный периоды.

Ключевые слова: дефицит витамина D, 25-гидроксивитамин D, общие группы риска, беременность, преждевременные роды.

The incidence of vitamin D deficiency in the population varies from 20% to 90% (2). The prevalence of vitamin D deficiency and insufficiency during pregnancy ranges from 19.0 to 96.0% in Europe, from 27.0 to 91.0% in the United States, from 39.0 to 65.0% in Canada, in Australia and New Zealand - from 25.0 to 87.0% and in Asia - from 45.0 to 100.0% (5). According to a meta-analysis of 195 vitamin D status studies conducted in 44 countries and involving more than 168,000 people, the average serum level of 25-hydroxyvitamin D (25(OH)D) in the population varies widely from 1.9 to 54.6 ng/ml. At the same time, 37.3% of researchers reported vitamin D deficiency (<20 ng/ml) in the population. It has been shown that vitamin D deficiency in the body of a pregnant woman is associated with an increased risk of premature birth, the development of placental insufficiency, preeclampsia, gestational diabetes mellitus, bacterial vaginosis, impaired contractile activity of the uterus, and an increase in the frequency of cesarean sections. In the fetus and newborn it leads to low birth weight, decreased femoral growth in utero, infant heart failure, craniotabes, acute lower respiratory tract infection, and hypocalcemia. Vitamin D deficiency is considered to be a negative factor affecting women's reproductive health (7,11).

The review is devoted to calcitriol-dependent mechanisms of the pathogenesis of such obstetric complications as spontaneous miscarriage, premature birth, fetal growth restriction, preeclampsia, perinatal infections and gestational diabetes mellitus. In light of the achievements of recent years, the role of vitamin D in the regulation of reproductive function has been proven. As a steroid hormone, this vitamin is necessary for a wide range of physiological processes. Vitamin D deficiency is currently a worldwide health problem and causes a very large percentage of acute and chronic pathologies. It actively influences various metabolic processes and takes part in the regulation of cell growth and functioning in the human body (6). It was reported that pregnant women with serum vitamin D concentrations less than 20 ng/mL were 3.81 times more likely to have preterm birth than women with vitamin D concentrations greater than 40 ng/mL (18).

Vitamin D is formed in the skin under the influence of ultraviolet radiation or comes from food, then a chain of metabolic processes occurs with the formation of active metabolites of vitamin D, which, together with parathyroid hormone and calcitonin, provide regulation of calcium and phosphate metabolism - the so-called

classical action of vitamin D. In pregnant women, syncytiotrophoblast is one of the sources of calcitriol production. During pregnancy, closer to 12 weeks, the concentration of 1,25(OH)₂D more than doubles, which confirms the important role of vitamin D in supporting gestation and controlling fetal development processes. It regulates decidualization, implantation, and expression of placental lactogen. In addition, vitamin D affects the secretion of progesterone and estriol, human chorionic gonadotropin (hCG), calcium absorption by the placenta and placental immunomodulation. Thus, active metabolites of vitamin D have been shown to affect numerous physiological processes. Currently, the term vitamin D combines more than 6 molecules of a similar nature, of which 2 molecules of steroid prohormones - D₂ and D₃ - have the greatest biological significance. Vitamins D₂ and D₃ are biologically inert. To convert them into the active form of D-hormone (1,25((OH))₂D), which binds to its receptors in tissues, two successive hydroxylations are necessary. The first occurs in the liver and is converted by the mitochondrial enzyme vitamin D 25-hydroxylase (CYP27A1) to 25-hydroxyvitamin D (25((OH))D), also known as calcidiol. The second hydroxylation occurs in the kidneys, resulting in the synthesis of biologically active 1,25-dihydroxyvitamin D (1,25(OH)₂D), or calcitriol, under the action of mitochondrial 1 α -hydroxylase (CYP27B1). The enzyme 1 α -hydroxylase is also found in many other tissues, which contributes to the local conversion of 25(OH)D to active Vitamin D levels influence 172 key physiological indicators of human health associated with the risk of developing various diseases and the vitamin D receptor (VDR) is found in more than 38 organs and tissues (20). There are “classical” effects of D-hormone associated with its influence on calcium-phosphorus metabolism and bone mineral density, and “non-classical” biological effects. The “non-classical” (extraosseous) effects of D-hormone include inhibition of cell proliferation and angiogenesis, stimulation of the production of insulin and cathelicidins (antimicrobial peptides), inhibition of renin production, anti-inflammatory, immunomodulatory, antibacterial, antitumor and a number of other properties.

Similar results were obtained in a study by some co-workers, which indicates that vitamin D in human ovarian granulosa cells in vitro stimulates the production of estradiol, estrone, progesterone, and insulin-like growth factor 1 (19). Reports on the prevalence of vitamin D deficiency and insufficiency in various countries around the world show the widespread prevalence of this problem among different population groups, regardless of the standard of living in the country and its geographical location. In the epidermis, cholecalciferol binds to vitamin D-binding protein and 70% of it comes from the bloodstream to the liver, and the other part goes to fat cells, where a vitamin D depot is formed. It has been shown that when human skin is exposed to sunlight at a single erythemal dose, the level of vitamin D₃ in the blood increases in the same way as after ingestion of 10,000 IU of vitamin D₃ (3). However, the

development of hypervitaminosis D during prolonged sun exposure does not occur due to blocking the entry of excess vitamin D from the skin into the bloodstream and its transformation into inactive compounds. With age, the content of 7-dehydrocholesterol in the epidermis decreases; accordingly, the synthesis of vitamin D₃ decreases and after 65 years its level decreases by more than 4 times. 25-hydroxycholecalciferol is considered the most accurate indicator of vitamin D levels. This is because 25(OH)D has a long half-life (about 3 weeks)(9).

The currently used 500 IU per day is sufficient to maintain the optimal level of metabolism of Ca and P, but is not sufficient to implement the non-calcemic functions of cholecalciferol. The main manifestations of the activity of its metabolites in the body are: regulation of mineral metabolism, especially calcium and phosphorus; regulation of bone growth, remodeling and repair; inhibition of the renin-angiotensin system; anti-inflammatory, regulation of cell growth and differentiation, angiogenesis; regulation of neuromuscular conduction (13). When prescribing vitamin D preparations, laboratory monitoring is recommended 2–3 months after the start of treatment. To assess vitamin D status in pregnant women, the total concentration of 25(OH)D [25(OH)D₂ and D₃] in blood serum was determined using an enzyme immunoassay method (7). The results of 25(OH)D levels were assessed in accordance with European recommendations. At a vitamin D level <20 ng/ml, its deficiency was diagnosed; at a concentration of 20–30 ng/ml – deficiency (suboptimal status); 30–100 ng/ml was accepted as the optimal level. To prevent vitamin D deficiency, pregnant and lactating women need to receive at least 800-1,200 IU daily. 90 pregnant women were observed, women received 5,000 units of vitamin D daily until 26 weeks of pregnancy, and the control group (44 patients) received placebo. As a result, women taking placebo had a statistically higher rate of development than those in the study group (35.9% vs. 10.9% $P < 0.005$), indicating the effectiveness of vitamin D supplementation in preventing preterm birth in the first and second trimesters of pregnancy (8,12,22). Summarizing the available data, it can be assumed that women at risk of preeclampsia, gestational diabetes mellitus, recurrent miscarriage, placental insufficiency and IUI should take at least 4000 IU of vitamin D at the preconception stage and from early pregnancy (14). For healthy women with a vitamin D level of 25-30 ng/ml, 2000 IU may be used; above 30 ng/ml, the currently recommended dose is 1000 IU (17,21). Preventing prematurity is a global priority; thus, in the future, women at risk of vitamin D deficiency may be assessed and vitamin D supplements prescribed.

Literature:

1. Абашова Е.И., Ярмолинская М.И., Мишарина Е.В. Роль витамина D в репродукции // Доктор.Ру. 2018. № 10 (154). С. 37–42.

2. Розикова, Д. К., & Ихтиярова, Г. А. (2023). THE STRUCTURE OF REPRODUCTIVE LOSSES IN UZBEK WOMEN. *ЖУРНАЛ РЕПРОДУКТИВНОГО ЗДОРОВЬЯ И УРО-НЕФРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ*, 4(4).
3. Салимова, Т. Б., & Дустова, Н. К. (2022). Роль доплерометрии при ранней диагностики синдрома отставания роста плода. *Новый день в медицине*, 8, 46.
4. Салимова, Т. Б., & Дустова, Н. (2023). ПРИЧИНЫ РИСКА РАЗВИТИЯ СИНДРОМА ОТСТАВАНИЯ РОСТА ПЛОДА В БУХАРСКОЙ ПОПУЛЯЦИИ. *ЖУРНАЛ РЕПРОДУКТИВНОГО ЗДОРОВЬЯ И УРО-НЕФРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ*, 4(4).
5. Хайдарова, Н. Б. (2023). Прогноз Недостаточности Плацентарной Системы У Беременных, Получавших Лечение От Covid-19 И Sars-Cov-2. *Central Asian Journal of Medical and Natural Science*, 4(5), 693-700.
6. Шарипова, Н. М. (2023). Влияние Дефицита Витамина D На Течение Беременности. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(10), 59-63.
7. Шарипова, Н. М. (2023). ВЛИЯНИЕ ДЕФИЦИТА ВИТАМИНА D НА ТЕЧЕНИЕ БЕРЕМЕННОСТИ ПРИ ПРЕЖДЕВРЕМЕННЫХ РОДАХ. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(11), 191-196.
8. Bahodirovna, H. N. (2023). Prognosis of Placental Insufficiency in Pregnant Women Treated for Coronavirus. *Eurasian Medical Research Periodical*, 20, 228-236.
9. Bahodirovna, H. N. (2023). COVID-19 VA SARS-CoV-2 DAN DAVOLANGAN HOMILADOR AYOLLARDA PLATSENTAR TIZIM YETISHMOVCHILIGINI BASHORAT QILISH. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(11), 235-241.
10. Bakhtiyarovna, S. T. (2023). The Reasons of Developing Intrauterine Growth Restriction Syndrome. *Central Asian Journal of Medical and Natural Science*, 4(5), 742-749.
11. Rozikova , D. K. (2023). THE IMPACT OF SUBCHORIONIC HEMATOMA ON THE FINAL RESULT OF PREGNANCIES IN INDIVIDUALS EXPERIENCING THREATENED ABORTION. *GOLDEN BRAIN*, 1(28), 57–62.
12. Rozikova Dildora Kodirovna. (2023). The Pattern of Reproductive Losese among Women in Uzbekistan’s Population. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(8), 52–60.
13. Salimova, T. B. (2022). Features of the Course of Pregnancy in Pregnant Women with Fetal Growth Restriction Syndrom and the Role of Doppler Velocimetry. *Central Asian Journal of Medical and Natural Science*, 3(6), 557-563.

14. SALIMOVA, T., & DO'STOVA, N. Q. (2023). HOMILA O'SISHINING CHEGARALANISHI SINDROMI BILAN HOMILADOR AYOLLARDA HOMILADORLIKNING KECHISHI XUSUSIYATLARI. *Молодые ученые*, 1(15), 4-6.
15. Salimova, T. (2023). CAUSES AND DIAGNOSIS OF INTRAUTERINE GROWTH RESTRICTION SYNDROME. *Science and innovation in the education system*, 2(11), 48-50.
16. Sharipova, N. M. (2023). Impact of Vitamin D Deficiency on Pregnancy. *Central Asian Journal of Medical and Natural Science*, 4(5), 705-712.
17. Sharipova, N. M. (2023). The Effect of Vitamin D Deficiency on The Course of Pregnancy During Premature Birth. *Central Asian Journal of Medical and Natural Science*, 4(6), 389-395.
18. T. B, S. . (2022). Homila O'sishi Chegaralanishi Sindromi Bilan Homilador Ayollarda-Homiladorlikning Kechishi Va Uning Diagnostikasida Dopplerometriyaning O'rni. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 1(6), 166–170. Retrieved from <https://www.sciencebox.uz/index.php/amaltibbiyot/article/view/4489>
19. Lanham-New S.A., Buttriss J.L., Miles L.M. et al. Proceedings of the rank forum on vitamin D. *Br J Nutr*. 2011;105(1):144–56.
20. Lisa M. Bodnar, Mark A. Klebanoff, Alison D. Gernand, Robert W. Platt, W. Tony Parks, Janet M. Catov, And Hyagriv N. Simhan Maternal vitamin D status and spontaneous preterm birth by placental Histology In The Us Collaborative Perinatal Project *Am J Epidemiol*. 2014;179(2):168–176
21. Farhan C, Eleftheria L, Katerina V, Guillermina G Immunomodulatory effects of vitamin D in pregnancy and beyond frontiers In *Immunology* November (2019) 10, 1-17
22. Kodirovna, R. D. (2023). The Effects of Subchorionic Hematoma on Pregnancy Outcome in Patients with Threatened Abortion. *Best Journal of Innovation in Science, Research and Development*, 2(10), 121–124.
23. Хамроев, X. H. (2022). Toxic liver damage in acute phase of ethanol intoxication and its experimental correction with chelate zinc compound. *European journal of modern medicine and practice*, 2, 2.
24. Khamroev, B. S. (2022). RESULTS OF TREATMENT OF PATIENTS WITH BLEEDING OF THE STOMACH AND 12 DUO FROM NON-STEROIDAL ANTI-INFLAMMATORY DRUGS-INDUCED OENP. *Journal of Pharmaceutical Negative Results*, 1901-1910.
25. Nutfilloyevich, K. K. (2023). STUDY OF NORMAL MORPHOMETRIC PARAMETERS OF THE LIVER. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(8), 302-305.

26. Nutfilloyevich, K. K. (2024). NORMAL MORPHOMETRIC PARAMETERS OF THE LIVER OF LABORATORY RATS. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(3), 104-113.

27. Nutfilloevich, K. K., & Akhrorovna, K. D. (2024). MORPHOLOGICAL CHANGES IN THE LIVER IN NORMAL AND CHRONIC ALCOHOL POISONING. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(3), 77-85.

28. Kayumova, G. M., & Hamroyev, X. N. (2023). SIGNIFICANCE OF THE FEMOFLOL TEST IN ASSESSING THE STATE OF VAGINAL MICROBIocenosis IN PRETERM VAGINAL DISCHARGE. *International Journal of Medical Sciences And Clinical Research*, 3(02), 58-63.

29. Хамроев, Х. Н., & Тухсанова, Н. Э. (2022). НОВЫЙ ДЕНЬ В МЕДИЦИНЕ. *НОВЫЙ ДЕНЬ В МЕДИЦИНЕ Учредители: Бухарский государственный медицинский институт, ООО "Новый день в медицине"*, (1), 233-239.

30. Хамроев, Х. Н. (2024). Провести оценку морфологических изменений печени в норме и особенностей характера ее изменений при хронической алкогольной интоксикации. *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ*, 36(3), 95-3.

31. Хамроев, Х. Н., & Тухсанова, Н. Э. (2021). Characteristic of morphometric parameters of internal organs in experimental chronic alcoholism. *Тиббиётда янги кун*, 2, 34.

32. Хамроев, Х. Н., Хасанова, Д. А., Ганжиев, Ф. Х., & Мусоев, Т. Я. (2023). Шошилинч тиббий ёрдам ташкил қилишнинг долзарб муаммолари: Политравма ва ўтқир юрак-қон томир касалликларига ёрдам кўрсатиш масалалари. *XVIII Республика илмий-амалий анжумани*, 12.

33. Хамроев, Х. Н., & Хасанова, Д. А. (2023). Жигар морфометрик кўрсаткичларининг меъёрда ва экспериментал сурункали алкоғолизмда қиёсий таснифи. *Медицинский журнал Узбекистана | Medical journal of Uzbekistan*, 2.

34. Khamroyev, X. N. (2022). TOXIC LIVER DAMAGE IN ACUTE PHASE OF ETHANOL INTOXICATION AND ITS EXPERIMENTAL CORRECTION WITH CHELATE ZINC COMPOUND. *European Journal of Modern Medicine and Practice*, 2(2), 12-16.

35. Hamroyev, X. N. (2022). The morphofunctional changes in internal organs during alcohol intoxication. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 2(2), 9-11.

36. Khamroyev, X. N. (2022). TOXIC LIVER DAMAGE IN ACUTE PHASE OF ETHANOL INTOXICATION AND ITS EXPERIMENTAL

CORRECTION WITH CHELATE ZINC COMPOUND. *European Journal of Modern Medicine and Practice*, 2(2), 12-16.

37. Хамроев, Х. Н. (2022). The morphofunctional changes in internal organs during alcohol intoxication. *EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE*, 2(2), 9-11.

38. Латипов, И. И., & Хамроев, Х. Н. (2023). Улучшение Результат Диагностики Ультразвуковой Допплерографии Синдрома Хронической Абдоминальной Ишемии. *Central Asian Journal of Medical and Natural Science*, 4(4), 522-525.

39. Хамроев, Х. Н., & Уроков, Ш. Т. (2019). ВЛИЯНИЕ ДИФФУЗНЫХ ЗАБОЛЕВАНИЙ ПЕЧЕНИ НА ТЕЧЕНИЕ И ПРОГНОЗ МЕХАНИЧЕСКОЙ ЖЕЛТУХИ. *Новый день в медицине*, (3), 275-278.

40. Хамроев, Х. Н., & Ганжиев, Ф. Х. (2023). Динамика структурно-функциональных нарушение печени крыс при экспериментальном алгоколние циррозе. *Pr oblems of modern surgery*, 6.

41. Уроков, Ш. Т., & Хамроев, Х. Н. (2018). Клинико-диагностические аспекты механической желтухи, сочетающейся с хроническими диффузными заболеваниями печени (обзор литературы). *Достижения науки и образования*, (12 (34)), 56-64.

42. Nutfilloevich, H. K., & Akhrorovna, K. D. (2023). COMPARATIVE CLASSIFICATION OF LIVER MORPHOMETRIC PARAMETERS IN THE LIVER AND IN EXPERIMENTAL CHRONIC ALCOHOLISM. *International Journal of Cognitive Neuroscience and Psychology*, 1(1), 23-29.

43. Ikhtiyarova, G. A., Dustova, N. K., & Qayumova, G. (2017). Diagnostic characteristics of pregnancy in women with antenatal fetal death. *European Journal of Research*, (5), 5.

44. Kayumova, G. M., & Nutfilloyevich, K. K. (2023). CAUSE OF PERINATAL LOSS WITH PREMATURE RUPTURE OF AMNIOTIC FLUID IN WOMEN WITH ANEMIA. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(11), 131-136.

45. Kayumova, G. M., & Dustova, N. K. (2023). Significance of the femoflor test in assessing the state of vaginal microbiocenosis in preterm vaginal discharge. Problems and scientific solutions. In *International conference: problems and scientific solutions. Abstracts of viii international scientific and practical conference* (Vol. 2, No. 2, pp. 150-153).

46. Каюмова, Г. М., Мухторова, Ю. М., & Хамроев, Х. Н. (2022). Определить особенности течения беременности и родов при дородовом излитии околоплодных вод. *Scientific and innovative therapy. Научный журнал по научный и инновационный терапии*, 58-59.

47. Kayumova, G. M., & Dustova, N. K. (2023). ASSESSMENT OF THE STATE OF THE GENITAL TRACT MICROBIOCENOSIS IN PREGNANT WOMEN WITH PREMATURE RUPTURE OF THE MEMBRANES USING THE FEMOFLOR TEST. *Modern Scientific Research International Scientific Journal*, 1(1), 70-72.

48. Valeryevna, S. L., Mukhtorovna, K. G., & Kobylovna, E. S. (2019). Premature Birth In A Modern Aspect. *International Journal of Bio-Science and Bio-Technology*, 11(10), 31-37.

49. Саркисова, Л. В., Каюмова, Г. М., & Умидова, Н. Н. (2018). Морфологические изменения фетоплацентарного комплекса при герпетической инфекции. *Тиббиётда янги кун*, 188-191.

50. Каюмова, Г. М., Саркисова, Л. В., & Умидова, Н. Н. (2018). Современные взгляды на проблему преждевременных родов. *Тиббиётда янги кун*, 183-185.

51. Каюмова, Г. М., Хамроев, Х. Н., & Ихтиярова, Г. А. (2021). Причины риска развития преждевременных родов в период пандемии организм и среда жизни к 207-летию со дня рождения Карла Францевича Рулье: сборник материалов IV-ой Международной научнопрактической конференции (Кемерово, 26 февраля 2021 г.). ISBN 978-5-8151-0158-6.139-148.

52. Саркисова, Л. В., Каюмова, Г. М., & Бафаева, Н. Т. (2019). Причины преждевременных родов и пути их решения. *Биология ва тиббиёт муаммолари*, 115(4), 2.

53. Kayumova, G. M., & Dustova, N. K. (2023). Significance of the femoflor test in assessing the state of vaginal microbiocenosis in preterm vaginal discharge. Problems and scientific solutions. In *International conference: problems and scientific solutions. Abstracts of viii international scientific and practical conference* (Vol. 2, No. 2, pp. 150-153).

54. KAYUMOVA, G., & DUSTOVA, N. (2023). Features of the hormonal background with premature surge of amniotic fluid. *Of the international scientific and practical conference of young scientists «Science and youth: conference on the quality of medical care and health literacy» Ministry of healthcare of the republic of kazakhstan kazakhstan's medical university «KSPH»*. ISBN 978-601-305-519-0.29-30.

55. Қаюмова, Г. М. НҚ Дўстова.(2023). Muddatdan oldin qog'onoq suvining ketishida xavf omillarning ta'sirini baholash. *Журнал гуманитарных и естественных наук*, 2(07), 11-18.

56. Каюмова, Г. М., & Мухторова, Ю. М. (2022). Пороговые значения антител к эстрадиолу, прогестерону и бензо [а] пирену как факторы риска преждевременного излития околоплодных вод при недоношенной

беременности. *Scientific and innovative therapy. Научный журнал по научный и инновационный терапии*, 59-60.

57. Sarkisova, L. V., & Kayumova, G. M. (2019). Exodius of premature birth. *Тиббиётда янги кун*, 1(25), 155-159.

58. Саркисова, Л. В., & Каюмова, Г. М. (2018). Перинатальный риск и исход преждевременных родов. *Проблемы медицины и биологии*, 169-175.

59. Каюмова, Г. М., Саркисова, Л. В., & Рахматуллаева, М. М. (2018). Особенности состояния плаценты при преждевременных родах. In *Республиканской научно практической конференции «Актуальные вопросы охраны здоровья матери и ребенка, достижения и перспективы»* (pp. 57-59).

60. Каюмова, Г. М., Саркисова, Л. В., & Саъдуллаева, Л. Э. (2018). Показатели центральной гемодинамики и маточно-фетоплацентарного кровотока при недонашивании беременности. In *Республиканской научно практической конференции «Актуальные вопросы охраны здоровья матери и ребенка, достижения и перспективы»* (pp. 56-57).

61. Саркисова, Л., Каюмова, Г., & Рузиева, Д. (2019). Современные тренды преждевременных родов. *Журнал вестник врача*, 1(4), 110-114.

62. Каюмова, Г. М., & Ихтиярова, Г. А. (2021). Причина перинатальных потер при преждевременных родов у женщин с анемией. (2021). In *Материалы республиканской научно-практической онлайн конференции. «Актуальные проблемы современной медицины в условиях эпидемии»* (pp. 76-7).

63. Kayumova, G. M., Khamroev, X. N., & Ixtiyarova, G. A. (2021). Morphological features of placental changes in preterm labor. *Тиббиётда янги кун*, 3(35/1), 104-107.

64. Khamroyev XN, Q. G. (2021). Improving the results of treatment of choledocholithiasis in liver diseases.

65. Kayumova, G. M. (2023). TO DETERMINE THE FEATURES OF THE COURSE OF PREGNANCY AND CHILDBIRTH IN WOMEN WITH PRENATAL RUPTURE OF AMNIOTIC FLUID. *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI*, 2(11), 137-144.

66. Kayumova, G. M. (2023). To Determine the Features Of Pregnancy and Children During Antenature Ruption Of Ambient Fluid. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 66-72.

67. Kayumova, G. M. (2023). Features of the Hormonal Background During Premature Relation of Ambitionall Fluid. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(9), 73-79.

68. Kayumova, G. M. (2023). The Significance Of Anti-Esterogen And Progesterone Antibodies As A Risk Factor In Premature Rupture Of Amniotic

Fluid. *American Journal of Pediatric Medicine and Health Sciences* (2993-2149), 1(9), 58-65.

69. Ro'ziyev, M. (2023, May). O'ZBEK VA NEMIS TILLARDA FE'LVNING MAJHUL NISBATINING IFODALANISHI. In *Integration Conference on Integration of Pragmalinguistics, Functional Translation Studies and Language Teaching Processes* (pp. 181-183).

70. Ro'ziyev, M. (2022). Periods of the educational process and the technology of teaching based on them. *ЦЕНТР НАУЧНЫХ ПУБЛИКАЦИЙ (buxdu.uz)*, 21(21).

71. Ro'ziyev, M. (2021). Ta'lim jarayonida yangi pedagogik texnologiyalardan foydalanishning psixologik imkoniyatlari. *ЦЕНТР НАУЧНЫХ ПУБЛИКАЦИЙ (buxdu.uz)*, 1(1).

72. Ro'ziyev, M. (2021). TA'LIM JARAYONINING UCHINCHI DAVRI ASOSIDA NEMIS TILI DARSLARINI O'TISH TEXNOLOGIYASI. *ЦЕНТР НАУЧНЫХ ПУБЛИКАЦИЙ (buxdu.uz)*, 3(3).

73. РЎЗИЕВ, М. К. ТАЪЛИМ ЖАРАЁНИДА ЯНГИ ПЕДАГОГИК ТЕХНОЛОГИЯЛАРДАН ФОЙДАЛАНИШНИНГ ПСИХОЛОГИК ИМКОНИЯТЛАРИ. *PSIXOLOGIYA Учредители: Бухарский государственный университет*, (S2), 50-52.

74. Ro'ziyev, M. (2020). Didaktik hodisalar tizimiga zamonaviy yondashuv. *ЦЕНТР НАУЧНЫХ ПУБЛИКАЦИЙ (buxdu.uz)*, 1(1).